

**Table 2. Estimated proportions of adults and adolescents with HIV/AIDS for whom a CD4 test was performed within 12 months after HIV diagnosis, by year of diagnosis and selected characteristics, 2001–2003—33 states with confidential name-based HIV infection reporting**

	2001			2002			2003		
	CD4 count <sup>a</sup>		Total No.	CD4 count <sup>a</sup>		Total No.	CD4 count <sup>a</sup>		Total No.
	No.	(%)		No.	(%)		No.	(%)	
<b>Sex</b>									
Male	15,734	(55.1)	28,555	15,895	(57.6)	27,617	16,083	(59.1)	27,200
Female	6,086	(49.7)	12,248	5,747	(51.0)	11,263	5,849	(54.8)	10,680
<b>Age as of end of year (yrs)</b>									
13–14	26	(48.4)	53	23	(50.2)	46	33	(61.7)	53
15–24	1,812	(42.7)	4,247	1,717	(41.2)	4,167	1,942	(44.5)	4,364
25–34	5,891	(51.9)	11,348	5,673	(52.6)	10,778	5,509	(55.3)	9,956
35–44	8,141	(54.5)	14,927	8,210	(58.3)	14,071	8,232	(60.7)	13,561
45–54	4,272	(57.6)	7,422	4,348	(61.2)	7,108	4,471	(61.9)	7,219
55–64	1,289	(60.4)	2,135	1,303	(62.5)	2,085	1,325	(62.9)	2,107
≥65	390	(58.2)	671	369	(58.9)	626	421	(67.8)	620
<b>Transmission category</b>									
<b>Male adult or adolescent</b>									
Male-to-male sexual contact	9,170	(55.3)	16,589	9,565	(56.9)	16,812	9,806	(58.5)	16,764
Injection drug use	2,865	(55.5)	5,162	2,689	(61.5)	4,370	2,577	(61.8)	4,169
Male-to-male sexual contact and injection drug use	916	(60.2)	1,522	921	(64.6)	1,427	888	(63.6)	1,396
Heterosexual contact <sup>b</sup>	2,662	(52.4)	5,081	2,600	(53.8)	4,829	2,695	(57.3)	4,702
Other <sup>c</sup>	121	(60.3)	200	120	(67.0)	179	117	(68.6)	170
<b>Female adult or adolescent</b>									
Injection drug use	1,440	(50.2)	2,869	1,290	(53.7)	2,403	1,277	(56.8)	2,250
Heterosexual contact <sup>b</sup>	4,527	(49.3)	9,173	4,347	(50.1)	8,678	4,450	(54.0)	8,236
Other <sup>c</sup>	119	(57.7)	207	110	(60.8)	181	121	(62.4)	195
<b>Race/ethnicity</b>									
White, not Hispanic	6,695	(59.9)	11,176	6,939	(61.4)	11,294	7,042	(63.8)	11,041
Black, not Hispanic	10,937	(51.4)	21,268	10,514	(52.6)	19,976	10,665	(55.8)	19,115
Hispanic	3,812	(50.0)	7,621	3,791	(55.0)	6,897	3,802	(54.1)	7,029
Asian/Pacific Islander	142	(53.8)	265	162	(52.1)	311	223	(61.4)	363
American Indian/Alaska Native	114	(66.8)	171	127	(63.6)	199	109	(58.3)	187
Unknown	120	(39.5)	303	109	(53.7)	202	91	(62.2)	146
<b>Total<sup>d</sup></b>	<b>21,820</b>	<b>(53.5)</b>	<b>40,804</b>	<b>21,642</b>	<b>(55.7)</b>	<b>38,880</b>	<b>21,932</b>	<b>(57.9)</b>	<b>37,880</b>

*Note.* These numbers do not represent actual cases in persons with a diagnosis of HIV infection or AIDS. Rather, these numbers are point estimates of case counts that have been adjusted for reporting delays and for redistribution of cases in persons initially reported without an identified risk factor. The estimates have not been adjusted for incomplete reporting.

Since 2000, the following states have had laws or regulations requiring confidential name-based HIV infection reporting: Alabama, Alaska, Arizona, Arkansas, Colorado, Florida, Idaho, Indiana, Iowa, Kansas, Louisiana, Michigan, Minnesota, Mississippi, Missouri, Nebraska, Nevada, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, South Carolina, South Dakota, Tennessee, Texas, Utah, Virginia, West Virginia, Wisconsin, and Wyoming. Since July 1997, Florida has had confidential name-based HIV infection reporting for new diagnoses only.

<sup>a</sup> When only CD4 percentage was available, a CD4 count was interpreted.

<sup>b</sup> With person at high risk or with a diagnosis of HIV infection or AIDS.

<sup>c</sup> Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or identified.

<sup>d</sup> Because column totals were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.